



# The routine childhood immunisation schedule for Wales

from November 2025

| Age due                                                     | Diseases protected against                                                                                    |                                                            | Vaccine and name                               |                            | Usual site <sup>1</sup> |
|-------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|------------------------------------------------|----------------------------|-------------------------|
| 8 weeks old                                                 | Diphtheria, tetanus, pertussis (whooping cough), polio, Haemophilus influenzae type b (Hib) and hepatitis B   |                                                            | DTaP/IPV/Hib/HepB (6-in-1)                     | Infanrix hexa® or Vaxelis® | Thigh                   |
|                                                             | Meningococcal group B (MenB)                                                                                  |                                                            | MenB                                           | Bexsero®                   | Thigh                   |
|                                                             | Rotavirus gastroenteritis                                                                                     |                                                            | Rotavirus                                      | Rotarix®                   | By mouth                |
| 12 weeks old                                                | Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B                                                    |                                                            | DTaP/IPV/Hib/HepB (6-in-1)                     | Infanrix hexa® or Vaxelis® | Thigh                   |
|                                                             | Meningococcal group B                                                                                         |                                                            | MenB                                           | Bexsero®                   | Thigh                   |
|                                                             | Rotavirus gastroenteritis                                                                                     |                                                            | Rotavirus                                      | Rotarix®                   | By mouth                |
| 16 weeks old                                                | Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B                                                    |                                                            | DTaP/IPV/Hib/HepB (6-in-1)                     | Infanrix hexa® or Vaxelis® | Thigh                   |
|                                                             | Pneumococcal                                                                                                  |                                                            | PCV13                                          | Prevenar 13® <sup>2</sup>  | Thigh                   |
| 12 months old                                               | <b>Born before 1 July 2024</b>                                                                                | <b>Born on or after 1 July 2024</b>                        |                                                |                            |                         |
|                                                             | Pneumococcal                                                                                                  | Pneumococcal                                               | PCV13                                          | Prevenar 13® <sup>2</sup>  | Upper arm/ thigh        |
|                                                             | Meningococcal group B                                                                                         | Meningococcal group B                                      | MenB                                           | Bexsero®                   | Thigh                   |
|                                                             | Measles, mumps and rubella                                                                                    | Measles, mumps and rubella                                 | MMR                                            | MMRVaxPro® or Priorix®     | Upper arm/ thigh        |
|                                                             | Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B                                                    |                                                            | DTaP/IPV/Hib/HepB (6-in-1)                     | Infanrix hexa® or Vaxelis® | Thigh                   |
| 18 months old                                               | <b>Born before 1 July 2024</b>                                                                                | <b>Born on or after 1 July 2024</b>                        |                                                |                            |                         |
|                                                             | No appointment                                                                                                | Measles, mumps and rubella                                 | MMR                                            | MMRVaxPro® or Priorix®     | Upper arm/ thigh        |
|                                                             |                                                                                                               | Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B | DTaP/IPV/Hib/ HepB (6-in-1)                    | Infanrix hexa® or Vaxelis® | Upper arm/ thigh        |
| 2 <sup>3</sup> and 3 years old and all school aged children | Influenza (annually from September)                                                                           |                                                            | Live attenuated influenza vaccine <sup>3</sup> | Fluenz® <sup>4</sup>       | Both nostrils           |
| 3 years 4 months old                                        | <b>Born before 1 July 2024</b>                                                                                | <b>Born on or after 1 July 2024</b>                        |                                                |                            |                         |
|                                                             | Diphtheria, tetanus, pertussis and polio                                                                      | Diphtheria, tetanus, pertussis and polio                   | dTaP/IPV (4-in-1)                              | Boostrix-IPV® or Repevax®  | Upper arm               |
|                                                             | Measles, mumps and rubella                                                                                    |                                                            | MMR                                            | MMRVaxPro® or Priorix®     | Upper arm               |
| School year 8 (12 to 13 year olds)                          | Cervical cancer, some head and neck and ano-genital cancers, and genital warts caused by human papillomavirus |                                                            | HPV <sup>5</sup> (one dose)                    | Gardasil 9®                | Upper arm               |
| School year 9 (13 and 14 year olds)                         | Tetanus, diphtheria and polio                                                                                 |                                                            | Td/IPV (3-in-1) (check MMR status)             | Revaxis®                   | Upper arm               |
|                                                             | Meningococcal groups A, C, W and Y                                                                            |                                                            | MenACWY                                        | Nimenrix® or MenQuadfi®    | Upper arm               |

<sup>1</sup> Where two or more injections are required at once, these should ideally be given in different limbs. Where this is not possible, injections in the same limb should be given 2.5cm apart. For more details see [Chapter 4 \(external site\)](#) and [Chapter 11 \(external site\)](#) in the [Green Book \(external site\)](#). All injected vaccines are given intramuscularly unless stated otherwise. Check relevant Green Book chapters regarding co-administration of vaccines.

<sup>2</sup> Refer to the [Green Book Chapter: Pneumococcal - table 25.3 \(external site\)](#) for babies with asplenia, splenic dysfunction, complement disorder or severely immunocompromised.

<sup>3</sup> Children must be 2 years old by 31 August to receive influenza vaccine in the routine programme in autumn/winter.

<sup>4</sup> If Fluenz® is contraindicated, use a suitable age-appropriate inactivated flu vaccine.

<sup>5</sup> Check the relevant chapter of the Green Book for individuals requiring a 3-dose schedule.

# Selective childhood immunisation programmes<sup>1</sup>

| Target group                                                                       | Age and schedule                      | Disease      | Vaccines                                                                                                                                                           |
|------------------------------------------------------------------------------------|---------------------------------------|--------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Babies born to hepatitis B infected mothers                                        | At birth and 1 month old <sup>2</sup> | Hepatitis B  | Hepatitis B vaccines<br>Engerix B® or HBvaxPRO®                                                                                                                    |
| Infants in areas of the country with TB incidence $\geq 40/100,000$                | From birth                            | Tuberculosis | BCG®                                                                                                                                                               |
| Infants with a parent or grandparent born in a high incidence country <sup>3</sup> | From birth                            | Tuberculosis | BCG®                                                                                                                                                               |
| Children in a risk group for influenza <sup>4</sup>                                | From 6 months to 17 years             | Influenza    | LAIV for children aged 2-17 years.<br>Appropriate inactivated flu vaccine for people aged 18 years and older or if LAIV is contraindicated or otherwise unsuitable |

<sup>1</sup> Other vaccines may be recommended for individuals with certain underlying medical conditions

<sup>2</sup> In addition to hexavalent vaccine (Infanrix hexa or Vaxelis) given at 8, 12, 16 weeks and 18 months. Take blood for HBsAg to exclude infection any time between 12 to 18 months.

<sup>3</sup> Where the annual incidence of TB is  $\geq 40/100,000$  see the [WHO country profile](#)

<sup>4</sup> Refer to annual flu Welsh Health Circular (WHC) for eligibility.



# Yr amserlen imiwneiddio reolaidd i blant ar gyfer Cymru o fis Tachwedd 2025

| Yr oedran i'w roi                                  | Yr afiechydon mae'n gwarchod rhag eu cael                                                          |                                                         | Brechiad ac enw                                |                             | Lleoliad arferol <sup>1</sup> |
|----------------------------------------------------|----------------------------------------------------------------------------------------------------|---------------------------------------------------------|------------------------------------------------|-----------------------------|-------------------------------|
| 8 wythnos oed                                      | Difftheria, tetanws, pertwsis (y pas), polio, Haemophilus influenzae math b (Hib) a hepatitis B    |                                                         | DTaP/IPV/Hib/HepB (6-mewn-1)                   | Infanrix hexa® neu Vaxelis® | Clun                          |
|                                                    | Grŵp meningococol B (MenB)                                                                         |                                                         | MenB                                           | Bexsero®                    | Clun                          |
|                                                    | Rotafeirws gastroenteritis                                                                         |                                                         | Rotafeirws                                     | Rotarix®                    | Drwy'r geg                    |
| 12 wythnos oed                                     | Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B                                            |                                                         | DTaP/IPV/Hib/HepB (6-mewn-1)                   | Infanrix hexa® neu Vaxelis® | Clun                          |
|                                                    | Grŵp meningococol B                                                                                |                                                         | MenB                                           | Bexero®                     | Clun                          |
|                                                    | Rotafeirws gastroenteritis                                                                         |                                                         | Rotafeirws                                     | Rotarix®                    | Drwy'r geg                    |
| 16 wythnos oed                                     | Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B                                            |                                                         | DTaP/IPV/Hib/HepB (6-mewn-1)                   | Infanrix hexa® neu Vaxelis® | Clun                          |
|                                                    | Niwmococol                                                                                         |                                                         | PCV13                                          | Prevenar 13 <sup>2</sup>    | Clun                          |
| 12 mis oed                                         | Ganwyd cyn 1 Gorffennaf 2024                                                                       | Ganwyd ar neu ar ôl 1 Gorffennaf 2024                   |                                                |                             |                               |
|                                                    | Niwmococol                                                                                         | Niwmococol                                              | PCV13                                          | Prevenar 13 <sup>2</sup>    | Rhan uchaf y fraich/clun      |
|                                                    | Grŵp meningococol B                                                                                | Grŵp meningococol B                                     | MenB                                           | Bexsero®                    | Clun                          |
|                                                    | Y frech goch, clwy'r pennau a rwbela                                                               | Y frech goch, clwy'r pennau a rwbela                    | MMR                                            | MMRVaxPro® neu Priorix®     | Rhan uchaf y fraich/ clun     |
|                                                    | Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B                                            |                                                         | DTaP/IPV/Hib/HepB (6-mewn-1)                   | Infanrix hexa® neu Vaxelis® | Clun                          |
| 18 mis oed                                         | Ganwyd cyn 1 Gorffennaf 2024                                                                       | Ganwyd ar neu ar ôl 1 Gorffennaf 2024                   |                                                |                             |                               |
|                                                    |                                                                                                    | Y frech goch, clwy'r pennau a rwbela                    | MMR                                            | MMRVaxPro® neu Priorix®     | Rhan uchaf y fraich/ clun     |
|                                                    | Dim apwyntiad                                                                                      | Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B | DTaP/IPV/Hib/ HepB (6-in-1)                    | Infanrix hexa® neu Vaxelis® | Rhan uchaf y fraich/ clun     |
| 2 <sup>3</sup> a 3 oed a phob plentyn oedran ysgol | Y Ffliw (bob blwyddyn o fis Medi)                                                                  |                                                         | Brechlyn fflw gwanedig byw (LAIV) <sup>3</sup> | Fluenz <sup>4</sup>         | Dau dwll y trwyn              |
| 3 blynedd a 4 mis oed                              | Ganwyd cyn 1 Gorffennaf 2024                                                                       | Ganwyd ar neu ar ôl 1 Gorffennaf 2024                   |                                                |                             |                               |
|                                                    | Difftheria, tetanws, pertwsis a pholio                                                             | Difftheria, tetanws, pertwsis a pholio                  | dTaP/IPV (4-mewn-1)                            | Boostrix-IPV® neu Repevax®  | Rhan uchaf y fraich           |
|                                                    | Y frech goch, clwy'r pennau a rwbela                                                               |                                                         | MMR                                            | MMRVaxPro® neu Priorix®     | Rhan uchaf y fraich           |
| Blwyddyn ysgol 8 (12 i 13 oed)                     | Canser serfigol, rhai canserau y pen a'r gwddw ac ano-genhedl, a dafadennau gwenerol a achosir gan |                                                         | HPV <sup>5</sup> (un dos)                      | Gardasil 9®                 | Rhan uchaf y fraich           |
| Blwyddyn ysgol 9 (13 a 14 oed)                     | Tetanws, difftheria a pholio                                                                       |                                                         | Td/IPV (3-mewn-1) (gwirio statws MMR)          | Revaxis®                    | Rhan uchaf y fraich           |
|                                                    | Grwpiau meningococol A, C, W ac Y                                                                  |                                                         | MenACWY                                        | Nimenrix® neu MenQuadfi®    | Rhan uchaf y fraich           |

<sup>1</sup> Pan fo angen dau bigiad neu fwy ar yr un pryd, dylid eu rhoi yn ddelfrydol mewn aelodau gwahanol. Os nad yw hyn yn bosibl, dylid rhoi pigiadau yn yr un aelod 2.5cm ar wahân. I gael rhagor o fanylion gweler [Pennod 4 \(safle allanol, Saesneg yn unig\)](#) a [Phennod 11 \(safle allanol, Saesneg yn unig\)](#) o'r [Llyfr Gwyrdd \(safle allanol, Saesneg yn unig\)](#). Rhoddir pob brechlyn i mewn i gyhyr, oni nodir yn wahanol. Gwiriwch benodau perthnasol yn y [Llyfr Gwyrdd](#) ynghylch cyd-weinyddu brechlynnau.

<sup>2</sup> Cyfeiriwch at [dabl 25.3 yn yr adran Niwmococol yn y Llyfr Gwyrdd \(safle allanol, Saesneg yn unig\)](#) sy'n trafod babanod ag asplenia, camweithrediad y ddueg, anhwylder sy'n effeithio ar y system ategol neu fabanod sydd ag imiwnedd gwan iawn.

<sup>3</sup> Rhaid i blant fod yn 2 oed erbyn 31 Awst i gael brechlyn fflw fel rhan o'r rhaglen reolaidd yn yr hydref/gaeaf.

<sup>4</sup> Os yw Fluenz® wedi'i wrtharwyddo, defnyddiwch frechlyn fflw wedi'i anactifadu sy'n briodol i'r oedran.

<sup>5</sup> Gwiriwch y bennod berthnasol yn y [Llyfr Gwyrdd](#) os oes gennych unigolion sydd angen amserlen 3 dos.

# Rhaglenni imiwneiddio dethol ar gyfer plant<sup>1</sup>

| Grwp targed                                                                                      | Oedran ac amserlen                      | Disease      | Vaccines                                                                                                                                                         |
|--------------------------------------------------------------------------------------------------|-----------------------------------------|--------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Babanod wedi'u geni i famau wedi'u heintio â hepatitis B                                         | Ar enedigaeth ac 1 mis oed <sup>2</sup> | Hepatitis B  | Brechiadau Hepatitis B<br>Engerix B® neu HBvaxPRO®                                                                                                               |
| Babanod mewn ardaloedd o'r wlad lle mae achosion o TB >= 40/100,000                              | O enedigaeth                            | Twbercwlosis | BCG®                                                                                                                                                             |
| Babanod gyda rhiant neu daid neu nain wedi'u geni mewn gwlad gyda llawer o achosion <sup>3</sup> | O enedigaeth                            | Twbercwlosis | BCG®                                                                                                                                                             |
| Plant mewn grŵp risg ar gyfer y ffliw <sup>4</sup>                                               | Rhwng 6 mis a 17 oed                    | Y ffliw      | LAIV ar gyfer plant 2-17 oed. Brechlyn ffliw wedi'i anactifadu priodol ar gyfer pobl 18 oed neu hŷn neu os yw LAIV wedi'i wrthgymeradwyo neu'n anaddas fel arall |

<sup>1</sup> Efallai y caiff brechiadau eraill eu hargymell ar gyfer unigolion gyda rhai cyflyrau

<sup>2</sup> Yn ogystal â brechlyn chwefalent (Infanrix hexa neu Vaxelis) a roddir yn 8, 12, 16 wythnos a 18 mis oed. Cymerwch waed i brofi am HBsAg er mwyn eithrio haint rhwng 12 a 18 mis oed.

<sup>3</sup> Pan fo nifer yr achosion blynyddol o TB yn >= 40/100,000 gweler [proffil gwlad Sefydliad Iechyd y Byd \(WHO\)](#)

<sup>4</sup> Cyfeiriwch at Gylchlythyr Iechyd Cymru (WHC) blynyddol ar y ffliw i gael manylion am gymhwysedd.